

## **JOB VI. POPULATION ASSESSMENT OF AMERICAN SHAD IN THE UPPER CHESAPEAKE BAY**

R.A. Sadzinski  
Maryland Department of Natural Resources  
Fisheries Service,  
301 Marine Academy Drive, Stevensville, MD 21666

### **INTRODUCTION**

The Maryland Department of Natural Resources has conducted annual sampling targeting adult American shad in the upper Chesapeake Bay since 1980 and hickory shad since 1998. The purpose of this sampling is to define stock characterizations including relative abundance indices, age and spawning history and reproductive success.

Since closure of the American shad fisheries to recreational and commercial fishing in 1980, stocks had increased significantly in the lower Susquehanna River until 2001 when American shad abundance decreased to alarming low levels. Hickory shad abundance appears to be very high and stable within the lower Susquehanna River. The Maryland Department of Natural Resources (DNR) is committed to restoring these species to the Susquehanna River Basin to sustainable, self-producing populations.

## METHODS AND MATERIALS

### I. Adults

#### a. American Shad

##### Field Operations

American shad were angled from the Conowingo tailrace (Figure 1) on the Susquehanna River two to five times per week from 27 April through 29 May 2009. Two rods were fished simultaneously, with each rod rigged with two shad darts and lead weight added, when necessary, to achieve proper depth.

All adult American shad sampled were sexed by expression of gonadal products (when possible) and fork length measured (mm). Scale samples were removed below the insertion of the dorsal fin. A minimum of three scales per fish were cleaned, mounted between two glass slides and read for age and spawning history using a Bell and Howell MT-609 microfiche reader. The scale edge was counted as a year-mark since it was assumed that each fish had completed a full year's growth at the time of capture. Fish in good physical condition and females not spent or running ripe were quickly tagged and released. A Maryland DNR Fisheries Service hat was given to fishers as reward for returned tags.

##### Statistical Analysis

The logistic growth model that described the population and catch estimate time-series was equivalent to a known biomass surplus production model (Macall 2002); except it used numbers as its unit of measure rather than biomass:

$$N_t = N_{t-1} \cdot (r \cdot N_{t-1} \cdot (1 - N_{t-1}) / K) - C_{t-1};$$

where  $N_t$  was the population in year  $t$ ;

$N_{t-1}$  was the population in the previous year;

$r$  was the intrinsic rate of population increase;

$K$  was the maximum population size; and

$C_{t-1}$  was losses associated with upstream and downstream fish passage in the previous year (equivalent to catch in a surplus production model).

We employed an observation error model that assumed all residual errors were in the population observations and the logistic equation used to describe the time-series was deterministic and without error (Haddon 2001). This model also assumed a proportional consumption of American shad by striped bass and also assumed that American shad were also landed as proportional bycatch to the Atlantic herring fishery because population estimates without these estimates fell below the catches at Conowingo Dam. Instantaneous annual fishing mortality rates associated with fish passage in year  $t$  ( $F_{tp}$ ) during 1984-2009 were estimated as  $\log_e[1 - (C_t / N_t)]$  (Ricker 1975). Estimates of abundance were derived from model output.

The Conowingo tailrace estimate utilized American shad captured in the tailrace and tagged and subsequently recaptured by the east fish lift. Fish caught in the east lift were dumped into a trough and directed past a 4'x10' counting window, identified to species, and enumerated by experienced Normandeau technicians. Hourly catch logs by species were subsequently produced and distributed to DNR personnel. American shad possessing a tag were counted and the tag color noted. Annual catch-per-unit-effort (CPUE) for American shad was calculated as the geometric mean of fish caught per lift hour. Annual CPUE of upper Bay American shad captured by hook and line was calculated as the geometric mean of fish caught per boat hour.

Data was also collected from two creel surveys targeting American shad in the lower Susquehanna River. One survey was a roving creel whereby tailrace anglers were visited on site and asked a series of questions regarding effort and success. The second survey required anglers to record their daily catch, location and hours fished in a logbook that was returned to the

Department at the end of the spring fishing season. For both surveys, CPUE was calculated as the number of fish caught per hour fished.

**b. Hickory Shad**

**Relative Abundance**

The annual CPUE of Deer Creek hickory shad was calculated as the number of fish caught-per-angler-hour and was obtained from spring logbook data returned from volunteer anglers.

**Mortality Estimates**

Two methods were utilized to estimate total instantaneous mortality of hickory shad based on scale repeat spawning marks. For the first method, hickory shad total instantaneous mortalities ( $Z$ ) were estimated by the  $\log_e$ -transformed spawning group frequency, plotted against the corresponding number of times spawned (assuming consecutive spawning; ASMFC 1988);

$$\log_e (S_{fx} + 1) = a + Z * W_{fx}$$

where  $S_{fx}$  = number of fish with 1,2,...f spawning marks in year x;

$a$  = y-intercept;

$W_{fx}$  = frequency of spawning marks (1,2,...f) in year x.

The second method averaged the differences between the natural logs of the spawning group frequency to provide an overall  $Z$  between age groups. The  $Z$  calculated for these fish represents mortality associated with repeat spawning.

## RESULTS

### **I. Adults**

#### **a. American shad**

##### **Sex and Age Composition**

The 2009 male-female ratio for Conowingo tailrace adult American shad captured by hook and line was 1.45:1. Of the 668 fish sampled by this gear, 622 were scale-aged (Table 1). Those American shad not aged directly because of regenerated scales, were not assigned ages.

Males were present in age groups 2-6 while females were found in age groups 4-9 (Table 1). The 2005 year-class of males (age IV) was the most abundant age group sampled, accounting for 66.1% of the total catch. For females, the 2005 (age IV) was the most abundant age group, accounting for 47.4%, of the total catch.

##### **Repeat Spawning**

The percentages of Conowingo tailrace repeat spawning American shad sampled by hook and line in 2009 was 20.3% for males and 18.2% for females (Table 1). The arcsine-transformed proportions of these upper Bay repeat spawners (sexes combined) has significantly increased for the time series ( $r^2 = 0.51$   $p < 0.001$ ; Figure 2).

##### **Relative Abundance**

During east lift operations from 1 April to 5 June 2009, clerks counted 29,272 American shad passing the viewing window. Peak passage was on 4 May when 4,670 American shad were recorded. Breakdown of the 151 marked fish observed is listed below.

| East Lift |             |                   |
|-----------|-------------|-------------------|
| Tag Color | Year Tagged | Number Recaptured |
| Orange    | 2009        | 104               |
| Green     | 2008        | 3                 |

| West Lift |             |                   |
|-----------|-------------|-------------------|
| Tag Color | Year Tagged | Number Recaptured |
| Orange    | 2009        | 44                |
| Green     | 2008        | 0                 |

In 2009, the west lift at Conowingo Dam operated from 28 April to 29 May. The 6,534 American shad caught in the west lift were returned to the tailrace, used for experimentation or retained for hatchery operations. Peak capture from the west lift was on 17 May when 1,069 American shad were recorded. Forty-four tagged American shad were recaptured in 2009 from the west lift

The Conowingo tailrace American shad relative population estimate in 2009 was 188,113 (Figure 3).

Estimates of hook and line and fish lift geometric mean (GM) CPUEs have increased significantly for the time series ( $r^2=0.31$ ,  $P= 0.004$ ;  $r^2=0.42$ ,  $P<0.001$ , respectively, Figures 4 and 5). However, a significant decrease in the fish lift GM CPUE ( $r^2=0.88$ ,  $P<0.001$ ) has been noted since 2002 while hook and line GM CPUE has declined in five of the previous seven years (Table 2).

The angler-based roving creel in the Conowingo Dam’s tailrace interviewed forty anglers in 2009 on only three days because of time constraints. Catch-per-angler-hour (CPAH) from these anglers was 1.41, a significant increase from 2008 when the CPAH was 0.74 (Table 3). CPAH from 2001-2009 has increased significantly ( $r^2=0.55$ ,  $P=0.02$ ). American shad logbook data indicated an increase in CPAH compared to 2008 (Table 4) but there was no significant trend for the time series (1999-2009,  $r^2=0.34$ ,  $P=0.06$ ).

**b. Hickory shad**

### Relative Abundance

Estimates of annual recreational hook and line (1998-2009) catch-per-angler-hour (CPAH) in Deer Creek ranged from 3.6 to 8.3 and have varied without trend since 1998 ( $r^2=0.14$ ,  $P=0.26$ ; Table 5).

### Mortality Estimates

Richardson (et al 2004) noted that ninety percent of hickory shad in Deer Creek have spawned by age four and this stock generally consisted of few virgin fish. The oldest fish in their sample was nine years old and using Hoenig's (1983) estimation of natural mortality ( $\ln(M_x) = 1.46 - 1.01 \{\ln(t_{\max})\}$ ),  $M$  was 0.47.

If  $Z$  is calculated using the freshwater spawning marks as in American shad, then mortality estimates for hickory shad estimated from the spawning group frequency plotted against the corresponding number of times spawned resulted in a  $Z$  of 0.59. The average difference between the natural logs of the spawning group frequency produced a  $Z$  of 0.54. In general, the resultant  $Z$  was attributed to natural mortality since both recreational and commercial fishing for hickory shad were banned.

## DISCUSSION

### Anadromous Species

#### 1. American shad

##### a. Adults

Prior to 1997, American shad captured from both fish lifts were individually handled so that all fish, both marked and unmarked, could be totaled. Beginning in 1997, the east fish lift became fully automated. Consequently, two trained observers stationed at the east lift-viewing window recorded both total counts and number of tagged American shad. This change in operating procedure at the east lift increased the chances of missing both tagged and untagged American shad and misidentifying tag colors. These errors could, therefore, affect the accuracy of the Petersen population estimates.

All American shad commercial fisheries in the Atlantic Ocean were closed on 31 December 2004. Since this fishery resulted in landings of mixed stocks in excess of 1.2 million lbs (ASMFC 1998) and a moratorium exists for American shad in the Chesapeake Bay, increases in relative abundances were expected. However, the three indicators of relative abundance (tailrace relative population estimates, hook and line geometric mean CPUEs, and Conowingo Dam lift geometric mean CPUEs) have shown declines since 2003.

Several factors contributing to this decline in abundance could be related to poor recruitment, Striped bass predation (Crecco et al 2006) and American shad harvested in the ocean as “bait”. Because of the difficulty in identifying and differentiating the four alosines, many subadults may be caught as bycatch, appearing as bait in various markets particularly in New England and southern Canada (K Hattala pers. comm, NY Dept. Env. Cons.).



## **b. Juveniles**

Baywide juvenile American shad production in 2009 decreased to very low levels for the time series (Figure 6). Although spawning conditions appeared favorable in the upper Chesapeake Bay in 2009, only two juvenile American shad were captured from the seven permanent seine sites and none from the six auxiliary stations. Juvenile American shad indices in the upper Chesapeake Bay (Figure 7) have been primarily driven by wild production below Conowingo Dam as indicated by the continued absence of hatchery-marked fish collected by the Juvenile Striped Bass Recruitment Survey (Project 2 Job 3 Task 3). This low juvenile index in 2009 for the upper Chesapeake Bay may demonstrate the decreasing trend in adult abundance and/or a change in environmental conditions

## **2. Hickory shad**

### **a. Adults**

Hickory shad are difficult to capture because of their aversion to fishery independent (fish lifts and traps) and fishery dependent (commercial pound and fyke nets) gears. Consequently, angler effort and success was collected from logbooks provided to anglers targeting hickory shad. Biological data and scale samples were obtained from hickory shad collected during electrofishing from Lapidum to the mouth of Deer Creek by DNR aquaculture personnel.

Deer Creek, a tributary to the Susquehanna River in Harford County has the greatest densities of hickory shad in Maryland (Richardson et al 2004). Natural mortality is approximately equal to the estimate of total mortality, demonstrating minimum mortality by hook and line and ocean bycatch.

### **b. Juveniles**

Haul seine sampling during the mid summer and fall likely missed hickory shad because of their large size, avoidance to the gear and their preference for deep water. Since adults may spawn from mid March through late April, up to six weeks before American shad, juvenile hickory shad reach a larger size earlier. Consequently, in order to accurately represent their juvenile abundance, sampling would need to be initiated by early June.

## Literature Cited

- ASMFC (Atlantic States Marine Fisheries Commission). 1988. Supplement to the fishery management plan for the anadromous alosid stocks of the eastern United States: American shad, hickory shad, alewife, and blueback herring. Washington, D. C.
- ASMFC (Atlantic States Marine Fisheries Commission). 1998. American shad stock assessment peer review report. Washington, D. C. 217 pp.
- Crecco, V. ,T. Savoy and J. Benway.2006. Stock Assessment of Connecticut River Shad: Examination of Fishing and Predation Effects on the Recent Stock Decline. Connecticut Marine Fisheries Division. Report to the American Shad Stock Assessment Subcommittee.
- Hoening, J. 1983. Empirical use of longevity data to estimate mortality rates. Fishery Bulletin 82(1):898-903.
- Jarzynski, A. R. Sadzinski. 2008. Stock Assessment of adult and juvenile anadromous species in the Chesapeake bay and select tributaries .Maryland Department Natural Resources, Report F-61-R-4 . Annapolis, Maryland.
- Leggett, W. C. 1976. The American shad *Alosa sapidissima* with special reference to its migration and population dynamics in the Connecticut River. American Fisheries Society Monograph 1:169-225.
- Richardson, B. M., R. P. Morin, M.W. Baldwin and C. P. Stence. 2004. Restoration of American shad and hickory shad in Maryland's Chesapeake Bay, Maryland Department of Natural Resources Annapolis, Maryland.
- Ricker, W. E. 1975. *Computation and interpretation of biological statistics of fish populations*. Fisheries Research Board of Canada Bulletin 191.

Table 1. Numbers of adult American shad and repeat spawners by sex and age sampled from the Conowingo tailrace in 2009.

Conowingo Dam Tailrace

| AGE             | Male  |         | Female |         | Total |         |
|-----------------|-------|---------|--------|---------|-------|---------|
|                 | N     | Repeats | N      | Repeats | N     | Repeats |
| 2               | 1     | 0       | 0      | --      | 1     | 0       |
| 3               | 20    | 0       | 0      | --      | 20    | 0       |
| 4               | 244   | 11      | 120    | 0       | 364   | 11      |
| 5               | 93    | 55      | 91     | 15      | 184   | 70      |
| 6               | 11    | 9       | 36     | 25      | 47    | 34      |
| 7               | 0     | --      | 5      | 5       | 5     | 5       |
| 8               | 0     | --      | 1      | 1       | 1     | 1       |
| 9               | 0     | --      | 0      | --      | 0     | --      |
| Totals          | 369   | 75      | 253    | 46      | 622   | 121     |
| Percent Repeats | 20.3% |         | 18.2%  |         | 19.4% |         |

Table 2. Conowingo Dam tailrace hook and line data, 1982-2009.

| Year | Total Catch | Hours fished | CPUE  | GM CPUE |
|------|-------------|--------------|-------|---------|
| 1982 | 88          | N/A          | N/A   | N/A     |
| 1983 | 11          | N/A          | N/A   | N/A     |
| 1984 | 126         | 52           | 2.42  | 1.07    |
| 1985 | 182         | 85           | 2.14  | 1.05    |
| 1986 | 437         | 147.5        | 2.96  | 1.85    |
| 1987 | 399         | 108.8        | 3.67  | 6.71    |
| 1988 | 256         | 43           | 5.95  | 6.54    |
| 1989 | 276         | 42.3         | 6.52  | 7.09    |
| 1990 | 309         | 61.8         | 5.00  | 3.6     |
| 1991 | 437         | 77           | 5.68  | 5.29    |
| 1992 | 383         | 62.75        | 6.10  | 5.05    |
| 1993 | 264         | 47.5         | 5.56  | 4.8     |
| 1994 | 498         | 88.5         | 5.63  | 5.22    |
| 1995 | 625         | 84.5         | 7.40  | 7.1     |
| 1996 | 446         | 44.25        | 10.08 | 9.39    |
| 1997 | 607         | 57.75        | 10.51 | 10.2    |
| 1998 | 337         | 23.75        | 14.19 | 9.86    |
| 1999 | 823         | 52           | 15.83 | 15.94   |
| 2000 | 730         | 35.75        | 20.42 | 13.98   |
| 2001 | 972         | 65.75        | 14.78 | 15.12   |
| 2002 | 812         | 60           | 13.53 | 15.94   |
| 2003 | 774         | 69.3         | 11.17 | 9.4     |
| 2004 | 474         | 38.75        | 12.23 | 9.48    |
| 2005 | 412         | 57.92        | 7.11  | 9.2     |
| 2006 | 360         | 33.75        | 10.28 | 7.61    |
| 2007 | 468         | 52.91        | 8.85  | 8.13    |
| 2008 | 164         | 39.85        | 4.12  | 3.14    |
| 2009 | 668         | 58.50        | 11.42 | 9.38    |

Table 3. Recreational creel survey data from the Susquehanna River below Conowingo Dam, 2001-2009.

| Year | Number of Interviews | Total Fishing Hours | Total Catch of American Shad | Mean Number of American shad caught per hour |
|------|----------------------|---------------------|------------------------------|--|
| 2001 | 90                   | 202.9               | 991                          | 4.88   |
| 2002 | 52                   | 85.3                | 291                          | 3.41   |
| 2003 | 65                   | 148.2               | 818                          | 5.52   |
| 2004 | 97                   | 193.3               | 233                          | 1.21   |
| 2005 | 29                   | 128.8               | 63                           | 0.49   |
| 2006 | 78                   | 227.3               | 305                          | 1.34   |
| 2007 | 30                   | 107.5               | 128                          | 1.19   |
| 2008 | 16                   | 32.5                | 24                           | 0.74   |
| 2009 | 40                   | 85.0                | 120                          | 1.41   |

Table 4. Summary of the spring American shad logbook data, 1999-2009.

| Year | Number of Returned Logbooks | Total Reported Angler Hours | Total Number of American Shad Caught | Mean Number of American Shad Caught Per Hour |
|------|-----------------------------|-----------------------------|--------------------------------------|--|
| 1999 | 7                           | 160.5                       | 463                                  | 2.88   |
| 2000 | 10                          | 404.0                       | 3137                                 | 7.76   |
| 2001 | 8                           | 272.5                       | 1647                                 | 6.04   |
| 2002 | 8                           | 331.5                       | 1799                                 | 5.43   |
| 2003 | 9                           | 530.0                       | 1222                                 | 2.31   |
| 2004 | 18                          | 750.0                       | 1035                                 | 1.38   |
| 2005 | 18                          | 567.0                       | 533                                  | 0.94   |
| 2006 | 19                          | 227.3                       | 305                                  | 1.34   |
| 2007 | 10                          | 285.5                       | 853                                  | 2.99   |
| 2008 | 16                          | 568.0                       | 1,269                                | 2.23   |
| 2009 | 10                          | 378                         | 967                                  | 2.60   |

Table 5. Summary of the spring hickory shad log book data from Deer Creek, 1998-2009.

| Year | Number of Returned Logbooks | Total Reported Angler Hours | Total Number of Hickory Shad Caught | Mean Number of Hickory Shad Caught per Hour |
|------|-----------------------------|-----------------------------|-------------------------------------|---|
| 1998 | 19                          | 600                         | 4980                                | 8.30  |
| 1999 | 15                          | 817                         | 5115                                | 6.26  |
| 2000 | 14                          | 655                         | 3171                                | 4.84  |
| 2001 | 13                          | 533                         | 2515                                | 4.72  |
| 2002 | 11                          | 476                         | 2433                                | 5.11  |
| 2003 | 14                          | 635                         | 3143                                | 4.95  |
| 2004 | 18                          | 750                         | 3225                                | 4.30  |
| 2005 | 18                          | 272.5                       | 1699                                | 6.23  |
| 2006 | 19                          | 762                         | 4905                                | 6.43  |
| 2007 | 17                          | 782.5                       | 3395                                | 4.34  |
| 2008 | 22                          | 995.25                      | 5,469                               | 5.50  |
| 2009 | 15                          | 561.25                      | 2,022                               | 3.60  |

Figure 1. Location of the hook and line sampling in Conowingo Dam tailrace in 2009.






Figure 2. Trends in arcsine-transformed percentages of repeat spawning American shad (sexes combined) collected from the Conowingo Dam tailrace (1984-2009).




Figure 3. Conowingo Dam tailrace relative estimates of American shad 1984-2009.




Figure 4. Geometric mean CPUEs from Conowingo Dam tailrace hook and line sampling, 1984-2009.




Figure 5. Geometric mean CPUE of American shad from the lifts at Conowingo Dam, 1980-2009.




Figure 6. Baywide juvenile American shad geometric mean CPUEs, 1959-2009.




Figure 7. Upper Chesapeake Bay juvenile American shad geometric mean CPUEs, 1959-2009.

